



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

LIST OF SALIENTIA FROM NEAR JACKSONVILLE, FLORIDA.

The following were collected ten miles south of Jacksonville by the writer, 1911-1912:

ORDER SALIENTIA.

Family *Ranidae*.

- Rana pipiens* Schreber, abundant all over.
Rana sphencephala Cope, abundant all over.
Rana æsopus Cope, local, not very plentiful.
Rana clamitans Latreille, sparsely, scattered companies.
Rana catesbiana Shaw, permanent bodies of water.
Rana gryllis Stejneger, not rare, but very shy.

Family *Engystomatidae*.

- Engystoma carolinense* Holbrook, abundant everywhere after heavy rains.

Family *Buфонidae*.

- Bufo lentiginosus* Shaw and
Bufo quercicus Holbrook, common, in companies in temporary water, or scattered on cultivated ground.

Family *Hylidae*.

- Hyla pickeringii* Storer, rare here.
Hyla squirella Bosc., in gardens and cornfields
Hyla femoralis Latreille, common on pine trees, but not easily caught.
Hyla cinerea Daudin, found in scattered bands, among water hyacinths on creeks.
Hyla gratiosa Le Conte, local, not very plentiful.
Chorophilus nigrinus Le Conte, common during the winter months.
Chorophilus occidentalis Bd. and Gd., plentiful, in scattered companies.
Chorophilus ocularis Holbrook, abundant everywhere.
Acris gryllus Le Conte, plentiful on edges of bayous.

R. F. DECKERT,
New York, N. Y.